

Schwarzschild or Kerr solution in 3-dim space and N -dim time

Dmitriy Palatnik *

August 1, 2009

Abstract

In this note Schwarzschild or Kerr solution is constructed in 3-dim space and N -dim time. Solutions do possess symmetry with respect to rotations in time volume.

1 Rotations in time plane

In paper [1] authors obtained spherically symmetric solution for metric for 2-dim time and 3-dim space in frames of Kaluza theoretic approach. For interval they obtained

$$dS^2 = dS_*^2 + \left(1 - \frac{J}{r}\right) dt^2 + \left(1 - \frac{C}{r}\right) du^2 + \frac{2P}{r} du dt \quad (1)$$

where

$$dS_*^2 = - \left(1 - \frac{R}{r}\right)^{-1} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (2)$$

Here u and t are time coordinates (speed of light $c = 1$) and R, J, C and P are constants satisfying relations

$$R = J + C ; \quad P^2 = JC \quad (3)$$

Substituting $P = \sqrt{JC}$ (where square root takes any of two signs) in (1) one obtains

$$dS^2 = dS_*^2 + dt^2 + du^2 - \frac{1}{r} \left(\sqrt{J} dt - \sqrt{C} du \right)^2 \quad (4)$$

Consider rotation in time plane

$$t = + \cos \tau \tilde{t} + \sin \tau \tilde{u} ; \quad (5)$$

$$u = - \sin \tau \tilde{t} + \cos \tau \tilde{u} \quad (6)$$

Metric (4) preserves its form under rotation (5) and (6) being provided with tilde quantities

$$\sqrt{\tilde{J}} = + \sqrt{J} \cos \tau + \sqrt{C} \sin \tau ; \quad (7)$$

$$\sqrt{\tilde{C}} = - \sqrt{J} \sin \tau + \sqrt{C} \cos \tau \quad (8)$$

Note that under rotation (7) (8)

$$\tilde{R} = \tilde{J} + \tilde{C} = J + C = R \quad (9)$$

From (3) it follows

$$\tilde{P} = P \cos 2\tau + \frac{1}{2}(C - J) \sin 2\tau \quad (10)$$

In particular taking

$$\tan \tau = \sqrt{\frac{C}{J}} \quad (11)$$

*The Waterford Nursing & Rehabilitation Centre 7445 N Sheridan Rd Chicago IL 60626-1818 USA

one obtains for interval

$$dS^2 = dS_*^2 + \left(1 - \frac{R}{r}\right) d\tilde{t}^2 + d\tilde{u}^2 \quad (12)$$

and in (12) one recognizes Schwarzschild solution in 4-dim spacetime with cylindrical extension on 2-nd time dimension which is also solution of Einstein equations in Ricci-flat 5-dim spacetime One may interpret worldline corresponding to $\{\tilde{u}; r; \theta; \phi\} = \text{constant}$ as one of uncharged black hole and rotation inverse to (5) (6) corresponds to situation when black hole has momentum along both time coordinates t and u which in Kaluza theory is adequate to motion of charged object In other words if one observer (frame of reference $\tilde{t}; \tilde{u}; r_o; \theta_o; \phi_o$) sees uncharged black hole then another observer (frame of reference $t; u; r_o; \theta_o; \phi_o$) sees same object as electrically charged These observations suggest insight for construction of analogous solution in case of N -dim of time volume

2 Schwarzschild solution for N -dim time

One may show that generalizing solution (1) on N -dim time one obtains following expression for interval

$$dS^2 = \left(\delta_{ik} - \frac{\sqrt{m_i m_k}}{r}\right) du^i du^k - \left(1 - \frac{M}{r}\right)^{-1} dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2) \quad (13)$$

Here $\{u^k\}$ $k = 1 \dots N$ are time coordinates; $\delta_{ik} = \text{diag}(1, 1, \dots, 1)$ is unit matrix $N \times N$; $\sqrt{m_k}$ ($= \pm \sqrt{m_k}$) are constants as well as M Really consider orthogonal transformation of time coordinates u^k ; $u_i = \delta_{ik} u^k$

$$u_i = \Omega_i^k \tilde{u}_k; \quad u^k = \Omega_i^k \tilde{u}^i \quad (14)$$

here

$$\Omega_i^k \Omega_l^i = \Omega_i^k \Omega_l^i = \delta_l^k \quad (15)$$

Taking

$$\sqrt{m_k} = \sqrt{M} \Omega_k^1 \quad (16)$$

and using (15) one obtains for interval

$$dS^2 = \left(1 - \frac{M}{r}\right) (d\tilde{u}^1)^2 + (d\tilde{u}^2)^2 + \dots + (d\tilde{u}^N)^2 - \left(1 - \frac{M}{r}\right)^{-1} dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2) \quad (17)$$

Evidently (17) is solution of Einstein equations hence (13) is solution as well

3 Kerr solution for N -dim time volume

Kerr solution in covariant and contravariant components [2] is

$$ds^2 = \left(1 - \frac{r_g r}{\rho^2}\right) dt^2 + \frac{2r_g r a}{\rho^2} \sin^2 \theta dt d\phi - \frac{\rho^2}{\Delta} dr^2 - \rho^2 d\theta^2 - \left(r^2 + a^2 + \frac{r_g r a^2}{\rho^2} \sin^2 \theta\right) \sin^2 \theta d\phi^2; \quad (18)$$

$$g^{ab} \partial_a \partial_b = \frac{1}{\Delta} \left(r^2 + a^2 + \frac{r_g r a^2}{\rho^2} \sin^2 \theta\right) (\partial_t)^2 + \frac{2r_g r a}{\rho^2 \Delta} \partial_t \partial_\phi - \frac{\Delta}{\rho^2} (\partial_r)^2 - \frac{1}{\rho^2} (\partial_\theta)^2 - \frac{1}{\Delta \sin^2 \theta} \left(1 - \frac{r_g r}{\rho^2}\right) (\partial_\phi)^2 \quad (19)$$

Here

$$\rho^2 = r^2 + a^2 \cos^2 \theta; \quad (20)$$

$$\Delta = r^2 + a^2 - r_g r \quad (21)$$

Take cylindrical extension of (18) on N -dim time $(\tilde{u}^1 \dots \tilde{u}^N)$ ($\tilde{u}^1 \equiv t$) which is also solution of Ricci-flat Einstein equations

$$dS^2 = \left(1 - \frac{r_g r}{\rho^2}\right) (d\tilde{u}^1)^2 + (d\tilde{u}^2)^2 + \dots + (d\tilde{u}^N)^2 + \frac{2r_g r a}{\rho^2} \sin^2 \theta d\tilde{u}^1 d\phi - \frac{\rho^2}{\Delta} dr^2 - \rho^2 d\theta^2 - \left(r^2 + a^2 + \frac{r_g r a^2}{\rho^2} \sin^2 \theta\right) \sin^2 \theta d\phi^2 \quad (22)$$

Consider rotation (14) (15) in time volume according to which $d\tilde{u}^l = \Omega_k^l du^k$ Then interval (22) may be rewritten as follows

$$dS^2 = \left(\delta_{kl} - \frac{r_g r}{\rho^2} \Omega_k^1 \Omega_l^1\right) du^k du^l + \frac{2r_g r a}{\rho^2} \Omega_k^1 \sin^2 \theta d\phi du^k - \frac{\rho^2}{\Delta} dr^2 - \rho^2 d\theta^2 - \left(r^2 + a^2 + \frac{r_g r a^2}{\rho^2} \sin^2 \theta\right) \sin^2 \theta d\phi^2 \quad (23)$$

Here $\delta_{kl} = \text{diag}(1, 1, \dots, 1)$ is unit matrix $N \times N$ Solution (23) corresponds to charged rotating black hole

4 Conclusion

In that work I have found easy way to generalize some of solutions of Einstein GRT on N -time volume

5 Acknowledgement

I wish to thank Christian network and Boris Tsirelson professor of Tel-Aviv University. I'm grateful for support from Catholic Church and Lubavich synagogue. I wish to apologize to Robert Geroch professor of University of Chicago for crimes I've execute in Chicago

References

- [1] J Kocinski M Wierzbicki arXiv: gr-qc/0110075
- [2] L D Landau E M Lifshits *The Classical Theory of Fields* Pergamon Press 1975